

VOL 2, No 54 (54) (2020)

The scientific heritage

(Budapest, Hungary)

The journal is registered and published in Hungary.

The journal publishes scientific studies, reports and reports about achievements in different scientific fields.

Journal is published in English, Hungarian, Polish, Russian, Ukrainian, German and French.

Articles are accepted each month.

Frequency: 24 issues per year.

Format - A4

ISSN 9215 — 0365

All articles are reviewed

Free access to the electronic version of journal

Edition of journal does not carry responsibility for the materials published in a journal.

Sending the article to the editorial the author confirms it's uniqueness and takes full responsibility for possible consequences for breaking copyright laws

Chief editor: Biro Krisztian Managing editor: Khavash Bernat

- Gridchina Olga Ph.D., Head of the Department of Industrial Management and Logistics (Moscow, Russian Federation)
- Singula Aleksandra Professor, Department of Organization and Management at the University of Zagreb (Zagreb, Croatia)
- Bogdanov Dmitrij Ph.D., candidate of pedagogical sciences, managing the laboratory (Kiev, Ukraine)
- Chukurov Valeriy Doctor of Biological Sciences, Head of the Department of Biochemistry of the Faculty of Physics, Mathematics and Natural Sciences (Minsk, Republic of Belarus)
- Torok Dezso Doctor of Chemistry, professor, Head of the Department of Organic Chemistry (Budapest, Hungary)
- Filipiak Pawel doctor of political sciences, pro-rector on a management by a property complex and to the public relations (Gdansk, Poland)
- Flater Karl Doctor of legal sciences, managing the department of theory and history of the state and legal (Koln, Germany)
- Yakushev Vasiliy Candidate of engineering sciences, associate professor of department of higher mathematics (Moscow, Russian Federation)
- Bence Orban Doctor of sociological sciences, professor of department of philosophy of religion and religious studies (Miskolc, Hungary)
- Feld Ella Doctor of historical sciences, managing the department of historical informatics, scientific leader of Center of economic history historical faculty (Dresden, Germany)
- Owczarek Zbigniew Doctor of philological sciences (Warsaw, Poland)
- Shashkov Oleg Candidate of economic sciences, associate professor of department (St. Petersburg, Russian Federation)

«The scientific heritage» Editorial board address: Budapest, Kossuth Lajos utca 84,1204 E-mail: public@tsh-journal.com Web: www.tsh-journal.com

CONTENT AGRICULTURAL SCIENCES

Dubrovskaya N.

EFFECTIVENESS OF FUNGICIDES AGAINST FUNGI FUSARIUM EQUISETI AND FUSARIUM LANGSETHIAE .3 *Rezvichkiy T., Tikidzhan R.,*

Mitlash A., Kalashnik V., Kochubey S.

BENEFIT OF SPRUCE. MAJOR DISEASES......5

Bekenev V.

THE RO	OLE C	DF N	ЛІТОС	HONDRIA	AND	FREE	RAD	ICAL
OXIDAT	ΓΙΟΝ	IN	THE	VIABILITY	OF	ANIM	۹LS	AND
HUMA	NS							13

Rezvichkiy T., Tikidzhan R.,

BIOLOGICAL SCIENCES

GEOLOGICAL AND MINERALOGICAL SCIENCES

Voronova E., Petrova L.

Voronova E., Petrova L.

MEDICAL SCIENCES

Kadiseva E., Kulagina L., Maksimov M., Ismailova M., Zvegintseva A.

VITAMIN THERAPY AND VITAMIN PREVENTION OF DISEASES OF THE MUSCULOSKELETAL SYSTEM27 *Abdurakhmanov I., Zhamilova G.,*

Yusupov A., Umurzakov Sh., Aidarov Z.

Murkamilov I., Sabirov I., Fomin V.,

Murkamilova Zh., Yusupov F.

 Tsoi L.

Pankov M., Afanasenkova N.,

Kuznetsova S., Cherska M.

VETERINARY SCIENCES

Serdyuc	henko l	l., Vasilyev	V., 1	Туи	tyunik A.
CHANGES	IN THE	INTESTINAL	MIC	ROF	LORA OF BEES
UNDER	THE	INFLUENCE		OF	ANTIBIOTIC
ENROFLOX	XACIN				

Serdyuchenko I., Vasilyev V., Tyutyunik A.

3. Долгих Л.А. Требования к составу и качеству рапсового масла для пищевой промышленности. VI международная конференция молодых ученых и специалистов, ВНИИМК, 2011 г. – С. 73–77. 4. Мхитарьянц Л.А. и др. Особенности химического состава семян рапса современных селекционных сортов // Масла и жиры. – Специализированный журнал. – 2016- № 3 (179).

INFLUENCE OF PLANT GROWTH REGULATORS ON ANATOMICAL OF FODDER BEAN LEAVES

Shevchuk O.

PhD of Biological Sciences, Associate Professor, Vinnytsia Mykhailo Kotsiubynskyi State Pedagogical University; Shevchuk V. Assistant Lecturer, Vinnytsia National Agrarian University

Abstract

It was studied the mesostructure organization of leaf apparatus and pigment content of leaves under application of growthregulators with different mechanism of action – esfon (2-HEFC) (0,2%) (ethylene producer) and Epin (0,1 ml/l) (growth stimulator) on broad bean plants. The plant growth regulators treatment on broad beans led to the thickening of leavesdue to increase in the growth of columnar and spongy leaf parenchyma. Growth regulators influenced on the formation of stomatal apparatus of broad beans leaves in different ways. The inhibitory compound esfon caused a decrease in the number of lower epidermal cells and stomata with a simultaneous increase in the stomatal cells area. The application of stimulator compound Epin led to an increase in epidermal cells, without differences in the number of stomata but the area of stomatal cells increased significantly. It has been established the enhancement of photosynthetic processes due to increase in the chlorophyll content in assimilative cells of leaves under esfon on broad beans.

Keywords: plant growth regulators, donor-acceptor system ("source-sink"), mesostructure organization, stomatal apparatus, chlorophyll, broad beans.

Formulation of the problem. The plant organism can be represented as the only donor-acceptor system ("source - sink"), in which the plant is considered as a system of donors (source) and acceptors (sink) of assimilates. The functioning of donor-acceptor system is determined by the genetic programof development. Donors and acceptors can be considered as individual plant structures - organs, tissues, cells and organelles, and as processes - photosynthesis, respiration, transport, assimilation (Kiriziy et al. 2014). It is known that regulation of source-sink system of a plant organism can be carried out artificially by growth regulators application, it causes an increase in crop production process (Kuryata & Khodanitska 2018; Shevchuk et al. 2019; Shevchuk 2020) and improve crop quality (Khodanitska et al. 2019; Shevchuk 2020) due redistribution of assimilate flow to economically valuable organs. Biologically active substances act on the regulatory mechanisms in cells of a plant organismat metabolic level (Bhatla 2018).

One of the modern method to increase the crop yield is an application of synthetic plant growth regulators and compounds that made on a natural substance basis and acts similarly to phytohormones (Kuryata et al. 2019; El Karamany 2019; Khodanitska et al. 2019). These compounds are environmentally friendly and positively affect on the microflora of soil, increase the resistance of plants to adverse environmental factors (extreme temperatures, drought and heat resistance, frost resistance) (Khan et al. 2019). The most common groups of synthetic plant growth regulators is retardants – antigibberellin compound. Retardants are widely used to increase the yield of oilseeds (Kuryata & Khodanitska 2018; Khodanitska et al. 2019; Kuryata et al. 2019), vegetables (Kuryata & Kravets 2018; Rohach et al. 2020), legumes (Kuryata et al. 2019; Shevchuk et al. 2020; Shevchuk 2020; Marchuk et al. 2016; Shevchuk et al. 2018; Shevchuk & Didur 2019) and industrial crops (Shevchuk et al. 2019; Shevchuk et al. 2017).

It is necessary to make attention to legumesdue to the growing deficiency of protein. The value of these crops lies in the fact that they not only increase the resources of food and feed grains but also increase soil fertility and crop production (Pankievicz et al. 2019). The broad bean plants havea great importance among legumes grown in our country for the grains and a green manure. They produce high yields on heavy clay soils compared to peas and fodder lupins, fix nitrogen from the atmosphere and is a good predecessor for winter and spring wheat. In this case, the issue of this study was to establish the application of stimulator Epin and inhibitor esfon (2-HEFC) on the mesostructure features of leaf apparatus of broad bean plants.

Conditions and methods of research. Broad bean plants cv. Vizir were treated with aqueous solutions of esfon (2-HEFC) (2 %) and growth stimulator Epin (0,1 ml/l) to complete wetting of leaves at shoot height 10-15 cm. Mesostructure organization of leaves were studied on a fixed materialaccording to the generally accepted method of Mokronosov A.T. and Borzenkova R.A.

It was used a mixture of equal parts of ethanol, glycerol and water with addition of 1 % formalin for preservation of leaves. The maceration agent was a 5 % solution of acetic acid in 2 mol/l hydrochloric acid. The method of partial maceration of leaf tissue was used to determine epidermal cells. Measurement of the area of epidermal cells was carried out by using a microscope and an ocular micrometer MOB-1-15x. The number of

cells in the tissue per unit area of the field of view was calculated with followed calculation of one cell and its volume. The chlorophyll content wasdetermined in fresh matter by spectrophotometric method on the spectrophotometer SF-16. The analytical repeatability of the research is fivefold (AOAS, 2010). The statistical processing of results was performed by the method of one-way analysis of variance using the computer program Microsoft Excel 2010.

Research results. The results of the study of anatomical structure of leaves indicate the restructuring of leaf apparatus of treated plants under application of growth inhibitor esfon and stimulator Epin on broad beans (Table 1). It was observed that the thickness of leaf bladeof treated plants increased due to the growth of parenchyma. It was found that both growth regulators increased the volume of columnar cells and the size (length and width) of spongy parenchyma cellsofleave. Table 1.

Leaf mesostructure indicators under plant growth i	regulators treatment	it on broad bean o	v. Vizir
Maggiramonta	Control	$E_{afon}(2.0\%)$	Epin $(0.1 m)$

Measurements	Control	Esfon (2 %)	Epin (0,1 ml/l)
Thickness of leave, µm	236,5±0,17	258,4±2,23*	262,1±7,48*
Volume of palisade parenchyma, μm^3	3989,6 ±70,03	4656,7 ± 52,61*	$6397,2 \pm 69,73*$
Length of spongy cells, µm	26,8±1,10	29,8±1,12	29,2±0,56
Width of spongy cells, µm	18,9±0,65	19,3±0,87	20,1±0,98
Number of stomatason 1 mm ² of the abaxial leaf surface, pieces	10,6±0,93	7,8±0,22*	11,6±0,28
Area of a stomata, μm ²	420,6±7=6,24	451,0±7,87*	479,9±5,87*
Number of epidermal cellson 1 mm ² of the abaxial leaf surface, pieces	27,7±0,37	18,4±0,42*	32,1±0,41

Note: * – difference is significant at p<0,05.

It was found an increase of columnar parenchyma that is the main assimilation tissue of leave. The stomata of lower epidermis are important for the transpiration processes – a physiological process that creates a continuous flow of water from the root system to leaves, combining all plant in a single system, and protects the plant organism from overheating.

Plant growth regulators influenced on the formation of stomatal apparatus of broad bean leaves. The drugs had different effects on the number of stomata. The number of stomata decreased under esfon treatment and their number remained practically unchanged under Epin. The application of inhibitory compound esfon led to decrease in the epidermal cells and the number of stomata with a simultaneous increase in the stomatal cell area. The treatment of stimulator Epin led to an increase in the number of epidermal cells, but the number of stomata changed and the area of stomatal cells increased significantly.

The effectiveness of photosynthesis is determined by the functional state of photosynthetic apparatus, one of the indicators of which is the pigment content. An increase in photosynthetic processes at the beginning of vegetation can be indicated by an increase in the chlorophyll content of broad bean plants cv. Vizir (Fig. 1). Application of 2 % esfon increased in the pigment content for 30 % compared to control. Growth stimulator Epin not affected on the pigment content, it was close to control at the last stages of vegetation.




Figure 1. Leaf chlorophyll content under growth regulators treatment on fodder bean, %. 1, 2, 3rd-5th, 10th, 15th day after treatment with growth regulators.

The analysis of mesostructural indicators of photosynthetic apparatus of broad bean plants confirmed that quaternary ammonium compound esfon (2 %) and growth stimulator Epin (0,1 ml/l) enhanced formation of photosynthetic apparatus: the number of stomata increased, the area of stoma and the thickness of leave due to the growth of chlorenchyma. It was noted that chlormequat chloridet reatment increased in photosynthetic processes due to thechlorophyllaccumulation in assimilative cells of broad bean leaves. It is advisable toincrease the crop production of broad bean plants by quaternary ethylene producer esfon (2 %) treatment.

Conclusion. Application of plant growth regulators is a powerful method regulate the activity of assimilative apparatus, as one of the main component of donor-acceptor system of plants, and can be used for targeted regulation of plastic substances redistribution in board beans, that increased in the number of stomata, the area of stoma, and the thickness of leave due to the growth of chlorenchyma. It testifies to the fact that esfon treatment on broad bean plants leads to an increase in the leaf chlorophyll content and improve photosynthetic processes.

References

1. AOAC. 2020. Official Meethods of Analysis of Association of Analytical Chemist International 18 th ed. Rev. 3. Asso of Analytical Chemist. Gaithersburg, Maryland, USA.

2. Bhatla, S.C. 2018. Plant Growth Regulators: An Overview. In: Plant Physiology, Development and Metabolism. Springe. Singapore. 559-568. https://doi.org/10.1007/978-981-13-2023-1_1

3. El Karamany, M.F., Sadak, M.S., Bakry, B.A. 2019. Improving quality and quantity of mungbean plant via foliar application of plant growth regulators in sandy soil conditions. Bull Natl Res Cent, 43. 61. https://doi.org/ 10.1186/s42269-019-0099-5

4. Kiriziy, D.A., Stasyk, O.O., Pryadkina, G.A., Shadchyna, T.M. 2014. Fotosintez. Assimilyatsiya CO2 i mehanizmy jejyo regulyatsii. Kiev.

5. Khan, N., Bano, A., Babar, M.A. 2019. The stimulatory effects of plant growth promoting rhizobacteria and plant growth regulators on wheat physiology grown in sandy soil. Arch Microbiol, 201. 769–785. https://doi.org/10.1007/s00203-019-01644-w

6. Khodanitska, O.O., Kuryata, V.G., Shevchuk, O.A. et al. 2019. Effect of treptolem on morphogenesis and productivity of linseed plants. Ukrainian Journal of Ecology. 9 (2). 119-126.

7. Khodanitska, O.O., Shevchuk, O.A., Tkachuk, O.O., Shevchuk, V.V. 2019. Features of the anatomical structure of the autonomic organs and flax oil yield (Linum usitatissimum L.) at applications growth stimulants. Science Rise: Biological Science. 4 (20). 35-40. DOI: 10.15587/2519-8025.2019.188317

8. Kuryata, V.G., Golunova, L.A., Poprotska, I.V., Khodanitska, O.O. 2019. Symbiotic nitrogen fixation of soybean-rhizobium complexes and productivity of soybean culture as affected by the retardant chlormequat chloride.Ukrainian Journal of Ecology. 9 (2). 5-13.

9. Kuryata, V.G., Khodanitska, O.O. 2018. Features of anatomical structure, formation and functioning of leaf apparatus and productivity of linseed under chlormequatchloride treatment. Ukrainian Journal of Ecology. 8 (1). 918-926. https://doi.org/10.15421/2018 294

10. Kuryata, V.G., Kravets, O.O. 2018. Features of morphogenesis, accumulation and redistribution of assimilate and nitrogen containing compounds in tomatoes under retardants treatment. Ukrainian Journal of Ecology. 8(1). 356-362.https://doi.org/10.15421/2018 222

11. Kuryata, V.G., Polyvanyi, S.V., Shevchuk, O.A. et al. 2019. Morphogenesis and the effectiveness of the production process of oil poppy under the complex action of retardant chlormequat chloride and growth stimulant treptolem. Ukrainian Journal of Ecology. 9 (1). 127-134.

12. Marchuk, Yu. M., Ilchenko, I.V, Matvienko, V.O, Biletska, I.V. et al. 2016. Influence of different plant growth regulators on seed productivity of fodder bean plants. Materials of the XII International Scientific-Practical Conference "Science Days – 2016". 16. 49–51.

13. Pankievicz, V.C.S., Irving, T.B., Maia, L.G.S. et al. 2019. Are we there yet? The long walk towards the development of efficient symbiotic associations between nitrogen-fixing bacteria and non-leguminous crops. BMC Biol. 17. 99. https://doi.org/10.1186/s12915-019-0710-0

14. Polyvanyi, S.V., Golunova, L.A., Baiurko, N.V., Khodanitska, O.O., Shevchuk, V.V., Rogach, T.I., Tkachuk, O.O., Knyazyuk, O.V., Zavalnyuk, O. L., Shevchuk O.A. 2020. Morphogenesis of mustard white under the action of the antigibberellic preparation chlormequat chloride. Modern Phytomorphology. 14. 101–103.

15. Rohach, V.V., Rohach, T.I., Kylivnyk, A.M., Polyvanyi, S.V., Bayurko, N.V., Nikitchenko, L.O., Tkachuk, O.O., Shevchuk, O.A. Hudzevych, L.S., Levchuk N.V. 2020. The influence of synthetic growth promoters on morphophysiological characteristics and biological productivity of potato culture. Modern Phytomorphology. 14. 111–114. https://doi.org/10.5281/zenodo.200121

16. Shevchuk, O.A., Kravets, O.O., Shevchuk, V.V., Khodanitska, O.O., Tkachuk, O.O. et al. 2020. Features of leaf mesostructure organization under plant growth regulators treatment on broad bean plants. Modern Phytomorphology. 14. 104–106. https://doi.org/10.5281/zenodo.200119

17. Shevchuk, O.A., Pervachuk, M.V., Vergelis, V.I. 2018. Influence of drugs of antihiberine action on germination of bean seeds. Bulletin of Uman National University of Horticulture. 1. 66–71. DOI 10.31395/2310-0478-2018-1-66-71

18. Shevchuk, O.A., Tkachuk, O.O., Kuryata, V.G., Khodanitska, O.O., Polyvanyi, S.V. 2019. Features of leaf photosynthetic apparatus of sugar beet under retardants treatment. Ukrainian Journal of Ecology. 9 (1). 115-120.

19. Shevchuk, V.V. 2020. Influence of stimulating drugs on qualitative characteristics of pea seeds of winter variety NS Moroz. Perspectives of world science and education Abstracts of VI International Scientific and Practical Conference. Osaka, Japan 26-28 February. 913–922.

20. Shevchuk, V.V. 2020. Symbiotic activity of peas under the action of microbial preparation and plant growth regulator. The 4th International scientific and practical conference "Actual trends of modern scientific research" (October 11-13, 2020) MDPC Publishing, Munich, Germany. 18–23.

21. Shevchuk, V.V. 2020. Comparative analysis of the influence of stimulant drugs on sowing characteristics of winter pea seeds and fodder beans. Dynamics of the development of world science. Abstracts of VII International Scientific and Practical Conference. Vancouver. Canada 18-20 March. 954–963.

22. Shevchuk, V.V., Didur, I.M. 2019. Influence of plant growth regulators on seedlings morphogenesis and laboratory seed germination of winter pea of variety NS Moroz. Bulletin of Uman National University of Horticulture. 2. 48–53. DOI 10.31395/2310-0478-2019-2-48-53 23. Shevchuk, V.V., Solodanyuk, Yu.V., Surzhik, V.V., Reyvakh, A.S., Stakh, V.V., Shevchuk, O.A. 2017. Indicators of the photosynthetic apparatus of sugar beet plants under retardation regulation. Modern scientific journal. 2(1). 27– 29.